

$SU(N)$ chiral gauge theories on the lattice: a quick overview

Maarten Golterman ^{a*} and Yigal Shamir ^b

^aDepartment of Physics and Astronomy, San Francisco State University, San Francisco, CA 94132, USA


^bSchool of Physics and Astronomy, Tel-Aviv University, Ramat Aviv, 69978 Israel

We describe how an $SU(N)$ chiral gauge theory can be put on the lattice using non-perturbative gauge fixing. In particular, we explain how the Gribov problem is dealt with. Our construction is local, avoids doublers, and weak-coupling perturbation theory applies at the critical point which defines the continuum limit of our lattice chiral gauge theory.

The construction of lattice chiral gauge theories (ChGTs) is an old problem. Because of the Nielsen–Ninomiya theorem [1] and the chiral anomaly [2], one either has to give up on chiral symmetry on the lattice, but enforce it to re-emerge in the continuum limit, or modify the lattice definition of chiral symmetry in order to maintain exact chiral symmetry on the lattice. For reviews, see refs. [3,4]. Here we follow the first approach, reporting on our recent completion of a construction based on non-perturbative gauge fixing [5]. We will have some comments on the second approach towards the end.

If gauge invariance is broken, the longitudinal modes of the gauge field couple to the fermions. If the dynamics of the longitudinal modes is uncontrolled, their “back-reaction” changes the fermion spectrum from chiral to vector-like [3]. A key point here is that having a reasonable definition of the fermion determinant for smooth gauge fields does not solve the problem.

A solution is to construct a lattice theory with a critical point whose universality class is described by the perturbative expansion of the target continuum theory [6]. A renormalizable gauge is mandatory [7], so that, in spite of the lack of gauge invariance of the regulated theory, we can use the usual power-counting arguments together with Slavnov–Taylor identities to construct the counter terms, which are finite in number.



Schematically, the critical point we are after is shown in the figure. g is the (bare) gauge coupling, and $1/\xi g^2$ is the parameter multiplying the gauge-fixing action. The vertical axis is the (relevant!) gauge field’s mass-squared counter term. (The axis corresponding to the bare coupling itself is not shown.) The usual lattice Yang–Mills critical point is at the origin. But here the longitudinal modes are uncontrolled, and destroy the chiral nature of the fermion spectrum. The critical point we aim for is point A , which involves both the limit $g \rightarrow 0$ (for example, at fixed ξ) and tuning the mass counter term to the critical line shown in the figure. (Obviously, all other counter terms need to be tuned as well, see below.)

In order to secure the desired critical point and recover the target chiral gauge theory, we need lattice perturbation theory (PT) to be valid near A . Whether this can be accomplished or not is a non-trivial non-perturbative question. As such, it cannot be addressed in the context of continuum perturbation theory, where it is simply

*Presenter at conference

assumed that some non-perturbative theory exists for which the usual perturbative expansion is valid. In a non-perturbative context one cannot assume the existence of this critical point, but instead one has to demonstrate that it exists.

Our actual construction of the appropriate critical point is achieved by using a lattice gauge-fixing action that 1) contains a lattice transcription of the longitudinal kinetic term $(\partial \cdot V)^2/2\xi g^2$ [7], and 2) has the unique minimum $U_\mu = \exp(iV_\mu) = 1$ [6]. This guarantees sufficient control over all short-distance effects (for details, see ref. [5]). In particular, no fermion doublers are generated if they are not present in the classical continuum limit [8,9].

If a regularization breaks (chiral) gauge symmetry, counter terms are needed to restore gauge invariance. One adjusts these counter terms such that the renormalized theory is invariant under BRST symmetry (this is possible to all orders for an anomaly-free fermion spectrum). In non-abelian theories, this raises the question as to how BRST symmetry works non-perturbatively. Our new work [5] answers this question.

We begin with reviewing the situation. One starts with $Z = \int dU \exp[-S_{inv}(U)]$, an un-fixed theory on the lattice with compact gauge fields. Following Faddeev–Popov, one inserts

$$Z_{gf} = \int d\phi d\bar{c} d\bar{c} db \exp[-S_{gf}(U^\phi, c, \bar{c}, b)], \quad (1)$$

$$S_{gf} = 2t[-i\text{tr}(bF(U)) + \text{tr}(\bar{c}\mathcal{M}c)] + \xi g^2 \text{tr}(b^2),$$

with $F(U)$ the gauge-fixing condition and $\mathcal{M}c = \delta_{BRST}F(U)$. The requirement is that Z_{gf} is independent of U , so that only a constant was inserted into Z . If this holds, the gauge-invariant correlation functions of the gauge-fixed theory are identical to those of the un-fixed theory.

The good news is that, thanks to BRST invariance, $dZ_{gf}/dt = 0$ [10]. Hence $Z_{gf}(1) = Z_{gf}(0)$ is indeed independent of U . The bad news is that $Z_{gf}(0) = 0$ [10] because, for $t = 0$, the Boltzmann weight is independent of the ghosts. The deeper reason for this zero is likely due to Gribov copies arising from $U(1)$ circles in the (compact) gauge group [11]. Obviously, if we cannot non-perturbatively gauge fix pure Yang–Mills, there is little hope of extending this to ChGTs.

Building on ref. [12], we proposed in ref. [5] to first fix only the coset $SU(N)/U(1)^{N-1}$ and then to fix the remaining $U(1)$ ’s without ghosts.

This procedure leads to a modified, “equivariant” BRST (eBRST) symmetry. Taking $SU(2)$ for example, we first split $V = \frac{1}{2}A\tau_3 + W^+\tau_+ + W^-\tau_-$, and choose the $U(1)$ -covariant gauge-fixing $F(V) = \partial \cdot W + i[A, W]$. We only introduce (anti-)ghosts $C^\pm$ ($\bar{C}^\pm$) for the generators $\tau_\pm$, which transform as

$$sC = (-iC^2)_{coset} \equiv -iC^2 + X, \quad s\bar{C} = -ib, \quad (2)$$

where also b lives only in the coset, and $X = iC_{U(1)}^2 \equiv (i/2)\tau_3\text{tr}(\tau_3C^2)$. We thus have $s^2C = -i[X, C] = \delta_X C$, a $U(1)$ transformation of C with parameter X . (For Ψ in the fundamental *irrep*, $s\Psi = -iC\Psi$ and $s^2\Psi = -iX\Psi$.) The equivariantly nil-potent algebra $s^2 = \delta_X$ requires $s^2\bar{C} = -i[X, \bar{C}]$, and therefore the new rule

$$sb = is^2\bar{C} = [X, \bar{C}] \neq 0. \quad (3)$$

eBRST is still nil-potent on any $U(1)$ -invariant operator. But, because of eq. (3), $\xi g^2 \text{tr}(b^2)$ is no longer eBRST invariant, and is replaced by

$$s(\xi g^2 \text{tr}(\bar{C}b)) = \xi g^2 [\text{tr}(b^2) + 4\bar{C}^+ C^+ \bar{C}^- C^-]. \quad (4)$$

The new 4-ghost term changes the bad into good news; we still have $dZ_{gf}/dt = 0$, but now

$$Z_{gf}(1) = Z_{gf}(0) = \int d\phi dC d\bar{C} db e^{-\xi g^2 [\text{tr}(b^2) + 4\bar{C}^+ C^+ \bar{C}^- C^-]} \neq 0, \quad (5)$$

is not equal to zero and independent of the gauge field, just as required. This is a rigorous result.

In order to construct a ChGT, complete gauge-fixing is needed. The remaining step is to fix the eBRST-invariant theory, which is now an abelian gauge theory, using a lattice version of $(\partial \cdot A)^2$ and no ghosts, in order to avoid the no-go theorem [10]. The resulting modified Slavnov–Taylor identities are still sufficient to prove unitarity (to all orders) [5]. The intuitive reason is that since the theories defined by $Z_{unfixed}$ and Z_{eBRST} have identical physical sectors, gauge fixing either one completely should not change this.

The lattice gauge-fixing action also contains an irrelevant term, which is needed for the uniqueness of the classical vacuum and, hence, for the

existence of a critical point near which PT is valid [6,5]. As a result, the fully gauge-fixed lattice Yang–Mills action is not eBRST invariant. However, since gauge invariance is already broken by the chiral fermions and counter terms are required anyway, this is not a new price to pay.

All these elements together lead to a complete construction of lattice $SU(N)$ ChGTs. Summarizing the main features, what we have is

- Locality,
- Perturbation theory near the critical point defining the continuum target theory,
- No fermion doublers, because of our control over the longitudinal modes,
- The theory accounts for fermion-number violation [13],
- Universality: our construction works for any lattice fermion method without doublers in the classical continuum limit [8,9].

What we do not have is

- Exact gauge invariance (we need a finite number of counter terms),
- Manifest unitarity; but it is recovered at least to all orders in perturbation theory.

In summary, our claim can be stated as follows: *If a certain chiral gauge theory exists, our construction provides a valid, non-perturbative lattice regularization.* For details, we refer to ref. [5].

In the remaining space we compare our results with those of refs. [14,15]. These references take the other route mentioned in the beginning: the modification of chiral symmetry on the lattice, with the aim of maintaining exact chiral gauge invariance. To do this, one employs Ginsparg–Wilson fermions, and left-handed fermions are defined through a modified chiral projection in which γ_5 is replaced by $\hat{\gamma}_5(U) = \gamma_5(1 - aD_{GW}(U))$. Because of the U dependence of $\hat{\gamma}_5$, the fermion measure is also U dependent. This leads to an integrability condition on the space of lattice gauge fields which has to be solved to define the theory [15]. The current state of affairs is that an exact solution was found for the

abelian case (using admissible gauge fields) [14]; for the non-abelian case, a solution was found only in perturbation theory [16] (requiring an infinite number of irrelevant counter terms), and no non-perturbative solution is known. (The Witten anomaly can be recovered [17].)

In the absence of a non-perturbative solution this approach is incomplete (see also ref. [4]). As it does not involve gauge fixing, there is no control over the dangerous “back reaction” of the longitudinal modes, unless gauge invariance is exact!

Our approach does not rely on exact gauge invariance and, as such, it does not “immediately” answer whether a certain unitary ChGT exists. But, whether it exists or not is a dynamical issue which *can* be investigated in our approach.

REFERENCES

1. H. B. Nielsen and M. Ninomiya, Nucl. Phys. B **193**, 173 (1981); Nucl. Phys. B **185**, 20 (1981) [Erratum-ibid. B **195**, 541 (1982)].
2. L. H. Karsten and J. Smit, Nucl. Phys. B **183**, 103 (1981).
3. D. N. Petcher, arXiv:hep-lat/9301015; Y. Shamir, arXiv:hep-lat/9509023.
4. M. Golterman, arXiv:hep-lat/0011027.
5. M. Golterman and Y. Shamir, arXiv:hep-lat/0404011.
6. Y. Shamir, Phys. Rev. D **57**, 132 (1998); M. Golterman and Y. Shamir, Phys. Lett. B **399**, 148 (1997).
7. A. Borrelli *et al.*, Nucl. Phys. B **333**, 335 (1990).
8. W. Bock, M. Golterman and Y. Shamir, Phys. Rev. Lett. **80**, 3444 (1998).
9. S. Basak and A. K. De, Phys. Rev. D **64**, 014504 (2001).
10. H. Neuberger, Phys. Lett. B **183**, 337 (1987).
11. M. Testa, Phys. Lett. B **429**, 349 (1998).
12. M. Schaden, Phys. Rev. D **59**, 014508 (1999).
13. M. Golterman and Y. Shamir, Phys. Rev. D **67**, 014501 (2003).
14. M. Lüscher, Nucl. Phys. B **549**, 295 (1999).
15. M. Lüscher, Nucl. Phys. B **568**, 162 (2000).
16. M. Lüscher, JHEP **0006**, 028 (2000).
17. O. Bär and I. Campos, Nucl. Phys. B **581**, 499 (2000).